

**Fig. 1. Serum homocysteine values.** (A) In 136 white patients on chronic dialysis (85 men, 51 women, age,  $61 \pm 14$  years) at baseline (before any vitamin supplementation) and 6 months after vitamin supplementation (vitamin B<sub>12</sub>, 1000 to 3000 μg/week plus folic acid 15 mg/week). Patients bearing the TT genotype ( $N = 22$ ; 16%) and never treated with vitamin therapies showed outstandingly higher serum homocysteine values as compared with patients with TC ( $N = 67$ ; 49%) and CC ( $N = 48$ ; 35%) genotypes. (B) Proportion of study patients by dialysis period in 136 white patients (85 males, 51 females, age  $61 \pm 14$  years) in whom the duration of dialysis was  $8.3 \pm 8.4$  years (minimum 0.6 year, maximum 33 years). Dialysis period is the year-subgroup of duration of dialysis. The percentage of TT homozygous in those with duration of dialysis of greater than 20 years was higher (18%) than in those with duration of dialysis <5 years (13%).

CATERINA CANAVESE, ALESSANDRA MESSUEROTTI,  
ROBERTA FENOGLIO, PAOLA MESIANO, PAOLA MASSARENTI,  
DANIELA PAULETTO, GIUSEPPE AIMO, AND GABRIELLA PRIOLO  
Torino, Italy

Correspondence to Dr. Caterina Canavese, Department of Nephrology S. Giovanni Molinette Hospital, Corso Bramante 88, 10126 Torino, Italy

E-mail: ccanavese@hotmail.com

## REFERENCES

- WRONE EM, ZEHNDER JL, HORNBERGER JM, *et al*: An MTHFR variant, homocysteine, and cardiovascular comorbidity in renal disease. *Kidney Int* 60(3):1106–1113, 2001
- BAGLEY SC, WHITE H, GOLOMB BA. Logistic regression in the medical literature: standards for use and reporting, with particular attention to one medical domain. *J Clin Epidemiol* 54(10):979–985, 2001
- ROSTAND SG, DRUEKE TB: Parathyroid hormone, vitamin D, and

cardiovascular disease in chronic renal failure. *Kidney Int* 56(2): 383–392, 1999

- BRATTSTROM L, ZHANG Y, HURTIG M, *et al*: A common methylenetetrahydrofolate reductase gene mutation and longevity. *Atherosclerosis* 141(2):315–319, 1998

## Reply from the Authors

Canavese *et al* were concerned (1) how to interpret the surprising association of lower homocysteine and cardiovascular comorbidity, (2) reconciling the difference in survival findings from our study and their earlier study, and (3) the cost benefit of genetic versus phenotypic screening.

The finding that higher homocysteine measured at baseline was associated with improved survival surprised us as well [1]. The epidemiologic findings of a risk factor having an apparently opposite association with disease in end-stage renal disease (ESRD) than it does in the

general population has been demonstrated for body size [2] and serum cholesterol [3] and has been postulated to be due to malnutrition [3]. We agree with Canavese et al that a great deal of research is still necessary regarding the biological mechanisms of homocysteine metabolism in patients with ESRD and their relationship to cardiovascular risk.

The finding of differing TT genotype prevalence over years of dialysis reported in these two studies is not irreconcilable. There may be other differences in the design of the studies that might explain variable risk estimates, including clinical status, sample size, measurement issues, and race/ethnicity composition. We intend to examine follow-up data from our study and agree that experience reported from other centers is greatly needed.

Routine screening of either homocysteine or genetics in not routinely recommended. A professional organization weighing the merits of such a clinical policy should take into consideration the cost-effectiveness of the test and associated interventions. In our clinic, the costs of such tests are significantly lower than those mentioned by Canavese et al.

ELIZABETH M. WRONE, JAMES L. ZEHNDER,  
JOHN M. HORNBERGER, LINDA M. McCANN,  
NORMAN S. COPLON, and STEPHEN P. FORTMANN  
Palo Alto, California, USA

Correspondence to Elizabeth Wrone, M.D., Stanford Center for Research in Disease Prevention, 1000 Welch Road, Palo Alto, California 94304-1825, USA.


E-mail: ewrone@stanford.edu

## REFERENCES

- WRONE EM, ZEHNDER JL, HORNBERGER JM, et al: An MTHFR variant, homocysteine, and cardiovascular comorbidity in renal disease. *Kidney Int* 60(3):1106–1113, 2001
- LEAVEY SF, STRAWDERMAN RL, JONES CA, et al: Simple nutritional indicators an independent predictors of mortality in hemodialysis patients. *Am J Kidney Dis* 31:997–1006, 1998
- CORESH J, LONGENECKER JC, MILLER ER, III, et al: Epidemiology of cardiovascular risk factors in chronic renal disease. *J Am Soc Nephrol* 9:S24–S30, 1998

## Daily hemodialysis and nutritional status

**To the Editor:** A switch from conventional hemodialysis (CHD) to daily hemodialysis (DHD) has been found to improve nutritional status of hemodialysis patients [1]. This observation by Galland et al needed to be confirmed. We, too, evaluated the effect of a switch from CHD (3 × 4 hours/week) to DHD (6 × 2 hours/week), using bioelectrical impedance analysis (BIA); we also observed a dramatic improvement in patients' lean body mass.



**Fig. 1. Time trends in evolution of lean body mass (LBM) after switch from conventional to daily hemodialysis in 7 patients with a 12-month follow-up.** It significantly increased within the first 6 months ( $P < 0.02$ ), and remained stable thereafter. Value at  $t_0$  represents the average between the values obtained at times  $-3$  and  $0$ , respectively. Mean weekly ( $\pm$  SEM) standard Kt/V [2] increased from  $2.22 \pm 0.13$  at baseline to  $3.25 \pm 0.20$  at 12 months ( $P < 0.001$ ).

Fourteen volunteer patients were evaluated during 3 months on CHD and 12 months on DHD. The lean body mass was evaluated at  $-3$ ,  $0$ ,  $+6$  and  $+12$  months, respectively. Five patients were excluded (renal transplantation in the first 6 months ( $N = 4$ ), incomplete data ( $N = 1$ ); data from the nine patients with 6 months DHD follow-up (six men, three women; 61 years old, range: 30 to 76 years), on CHD (five at home, three in a low-care unit, one in hospital) for 63 months (range: 8 to 137 months) was analysed by ANOVA for repeated measures with F or Friedman tests, as appropriate.

The lean body mass (mean  $\pm$  SEM) increased ( $+1.3 \pm 0.4$  kg) during the first 6 months ( $P < 0.008$ ) of DHD, while total body weight remained unchanged. After 6 months, two more patients were withdrawn (renal transplantation and aortic valve replacement, respectively). The lean body mass evolution of the remaining seven patients is presented (Fig. 1).

Based on these observations, a switch to DHD could be proposed to CHD patients with poor nutritional status. Further studies, including a larger number of patients, are required to confirm these encouraging results.

ERIC GOFFIN, YOLANDE PIRARD, JULIE FRANCART,  
MERCEDES VIGNIOBLE, TONY GOOVAERTS,  
ANNIE ROBERT, and YVES PIRSON  
Brussels, Belgium

## ACKNOWLEDGMENTS

The authors thank Dr. Ingrid Ledebro from Gambro, Sweden, for her support, and Professor M. Jadoul for his comments.

Correspondence to Eric Goffin, M.D., Department of Nephrology, Université catholique de Louvain, Cliniques Universitaires Saint-Luc, Av Hippocrate 10, 1200 Bruxelles, Belgium.  
E-mail: goffin@nefr.ucl.ac.be